

# Environmental sustainability in dermatological surgery. Part 1: reducing carbon intensity

Fatima Ali<sup>1,2</sup>, BDS Sustainability Subgroup Collaborative, Aaron Wernham<sup>3</sup>  
and Rachel Abbott<sup>4</sup>

<sup>1</sup>King's College London, London, UK

<sup>2</sup>Guy's and St Thomas' NHS Foundation Trust, London, UK

<sup>3</sup>Walsall Healthcare NHS Trust, Walsall, UK

<sup>4</sup>Cardiff and Vale University Health Board, Cardiff, UK

A full list of members of the BDS Sustainability Subgroup Collaborative is provided in [Appendix 1](#), and a list with affiliations in [Appendix S1](#) (see [Supporting Information](#)).

Correspondence: Fatima Ali. Email: [fatima.1.ali@kcl.ac.uk](mailto:fatima.1.ali@kcl.ac.uk)

## Abstract

This two-part review addresses the pressing need for environmental sustainability in dermatological surgery, driven by the National Health Service's commitment to net-zero emissions. Part 1 focuses on strategies to reduce the carbon intensity of dermatological procedures by adopting low-carbon alternatives and optimizing operational resource usage. Key strategies for a system-wide reduction in environmental impact include leveraging local suppliers to reduce transport emissions, streamlining care models, promoting efficient waste management and using mindful prescribing practices. Another aspect is integrating sustainability into dermatological education while minimizing the carbon footprint of surgical education. Additionally, the review provides a comprehensive overview of optimizing resource use in dermatological surgery, focusing on efficient management of consumables, equipment and energy. This includes recycling, waste segregation, transitioning to reusable personal protective equipment and surgical instruments, and applying energy-saving and sustainable water use practices. By implementing these strategies, dermatological surgery can significantly reduce its environmental impact while upholding high standards of patient care.

## Introduction

Growing environmental concerns have prompted a heightened focus on sustainability in the healthcare domain and, in particular, dermatological surgery. Climate change and dermatological surgery are (through complex and indirect pathways) causally connected, such that the delivery of the service itself is being affected by the unfolding crisis. This manifests in increased surgical rates due to a rise in skin pathologies related to climate change. The principles of planetary health guide our urgency to mitigate climate change and benefit the broader community served by dermatologists.

Sustainability involves meeting present requirements without compromising those of the future.<sup>1</sup> A sustainable dermatological surgical service delivers high-quality care while minimizing adverse environmental, social and economic impacts. In consequence, these services will adhere to the triple bottom line principle.<sup>2</sup> The National Health Service (NHS) is a major contributor to greenhouse gas emissions within the UK public sector and has committed to achieving net-zero emissions by 2045.<sup>3</sup>

Due to its resource-intensive nature, dermatological surgery generates a substantial carbon footprint.<sup>4</sup> This is further

compounded by the escalating incidence of skin cancer. It is estimated that over 200 000 surgical excisions are carried out in UK dermatology services each year, with over 170 000 new skin cancer diagnoses each year.<sup>5,6</sup> Dermatologists acknowledge this dual impact of climate change on skin health and the broader environment.<sup>7,8</sup>

Sustainability efforts in dermatological surgery encompass the consumption patterns in service provision, the prevention of skin diseases, streamlining of care models, clinical decision making, and broader collaborative partnerships within the healthcare pathway. The Centre for Sustainable Healthcare has outlined key principles to improve sustainability by reducing activity, reducing carbon intensity through opting for low-carbon alternatives, and being cognizant of resource use, research and innovation.<sup>9</sup>

This review aims to identify effective strategies and practices that can reduce the carbon footprint of dermatological surgical services while maintaining high standards of patient care. The review consists of two parts. Part 1 discusses strategies to reduce the carbon intensity of skin surgery through electing for low-carbon alternatives and environmentally optimizing day-to-day operational resource use. Part 2 evaluates methods to reduce the overall activity of

Accepted: 6 October 2024

© The Author(s) 2024. Published by Oxford University Press on behalf of British Association of Dermatologists. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

dermatological surgery and future environmental considerations within management, research and innovation.

## Methods

These articles aim to formally publish the original British Society for Dermatological Surgery Sustainability Guidance 2022<sup>1</sup> as narrative reviews with an updated literature search (to 30 June 2024). Details of separate phases of the literature scoping, theme identification, analysis and mapping to the Centre for Sustainable Healthcare key principles can be found in Part 2 of this review. Figure 1 is a summary infographic from the British Society for Dermatological Surgery (BSDS) Sustainability Guidance 2022.<sup>1</sup>

## Reducing carbon intensity in dermatological surgery


### Low-carbon alternatives

Low-carbon alternatives in dermatological surgery relate to treatment options and medical technologies with lower environmental impacts. The procurement process is carbon intensive and contributes to 22% of the NHS carbon footprint.<sup>10</sup> Dermatological surgery requires essential pharmaceutical agents such as local anaesthetics. However, it is important to consider whether there is a good indication for prescribing additional agents such as tranexamic acid and oral antibiotics.<sup>11</sup> A low-carbon alternative approach aims to decarbonize emissions-intensive areas of the supply chain, such as by using local suppliers to reduce transport-related carbon emissions. However, challenges arise due to increased legislation favouring mass manufacturing over local compounding.<sup>12</sup>

As the carbon emissions from procurement are directly correlated to supply chain demand, it is crucial to critically evaluate overtreatment practices. This involves determining whether specific interventions provide substantial benefits to patient outcomes and carefully prescribing the quantity and duration of treatments.<sup>13</sup> Such mindful prescribing practices can significantly mitigate pharmaceutical waste, which has an estimated monthly environmental impact of 15 kg CO<sub>2</sub> equivalent (CO<sub>2</sub>e) in wasted prescriptions (akin to an average car travelling 88 km).<sup>14</sup>

Pharmacists play a vital role in efficient dispensing, improving pharmaceutical preparation procedures, streamlining dispensing processes through electronic prescriptions and minimizing waste through effective stock management.<sup>15</sup> This approach requires empowering patients through awareness of carbon footprint labelling and providing transparent information on a product's environmental impact, shelf life and recycling options.<sup>16</sup> Dermatological surgery departments, the broader NHS and legislative bodies should advocate for environmental labelling on pharmaceuticals and skincare products.<sup>17,18</sup>

Road travel associated with NHS activities significantly contributes to England's total emissions, accounting for 3.5% of all road travel and 14% of the NHS system's emissions.<sup>19</sup> This encompasses business travel, fleet transport, patient travel, staff commutes and visitor travel. The NHS



**Figure 1** Summary infographic from the BSDS Sustainability Guidance 2022,<sup>1</sup> highlighting areas covered within this literature review. LA, local anaesthetic; LED, light-emitting diode; MDT, multidisciplinary team; MMS, Mohs micrographic surgery; NHS, National Health Service; PPE, personal protective equipment; QIP, quality improvement project; SSC, student-selected component. Figure reproduced with permission from the copyright holders.

transport fleet alone generates approximately 1000 kt CO<sub>2</sub>e annually.<sup>20</sup> Transitioning from car usage to low-carbon alternatives such as public transport, cycling or walking for both the skin surgery team and patients could potentially reduce

emissions by 461 kt CO<sub>2</sub>e per year.<sup>20</sup> Strategies for implementation include car-sharing initiatives, bicycle parking facilities, reward schemes, travel reimbursements and support for remote work.

Additionally, adopting care models such as ‘spot’ clinics, where consultant dermatologists visit community sites, can significantly decrease the need for patient travel.<sup>21,22</sup> However, implementing strategies that involve changes to service models poses several challenges, including initial infrastructure costs, logistical complexities, need for robust support systems, and careful planning to ensure equitable access and efficient scheduling.

A low-carbon alternative approach in dermatological surgery also integrates sustainability principles into dermatological education, which fosters awareness and implementation of eco-friendly practices. While the General Medical Council emphasizes sustainability education for newly qualified UK doctors, currently undergraduate and postgraduate education pay limited attention to climate change in curricula.<sup>23,24</sup> Surgical education is also resource intensive as it often uses single-use materials akin to healthcare delivery, and plastic manufacturing models for simulation training. Methods suggested to improve this include reducing dependence on single-use materials, use of materials aligned with circular economy principles, and low-carbon alternatives such as immersive augmented or virtual technologies.<sup>25</sup> More research is needed as current technologies do not accurately replicate organic tissues or simulate real surgical scenarios, which create unrealistic training environments.<sup>26</sup>

Medical conferences also significantly contribute to medical education’s carbon footprint through travel, energy consumption and food waste. Strategies to mitigate this impact include carbon-neutral conferences, virtual and hybrid formats, and reducing the carbon footprint of food.<sup>27</sup>

## Optimizing operational resource use in dermatological surgery

### Consumables

Consumables in dermatological surgery (equipment, packaging and personal protective equipment) contribute significantly to carbon emissions (Figure 2). Reducing unnecessary procedures can diminish use of all three types of consumables, as stated in the national specialty report for the Dermatology Getting It Right First Time programme.<sup>5</sup> Significant variation exists among UK providers regarding biopsies prior to definitive treatment of skin cancers, ranging from < 10% to over one-third of lesions. This variation causes greater resource use, excess waste and patient travel.<sup>5</sup> Strategies to mitigate this include avoiding unnecessary procedures such as biopsies, employing less resource-intensive approaches like topical therapies, and upskilling primary care providers to follow local referral pathways to reduce inappropriate referrals.<sup>28</sup>


### Equipment

#### Life-cycle assessment

Life-cycle assessment is a ‘cradle-to-grave’ analysis that estimates the environmental impact of different equipment used in skin surgery. This includes raw material extraction, manufacturing and transport.<sup>29</sup> Evaluating raw material utilization and manufacturing locations is essential in determining the carbon footprint.<sup>30</sup>

#### Single-use vs. multiuse surgical instruments

Historically, single-use surgical instruments gained preference due to perceived advantages in sterility and sharpness.<sup>31,32</sup>



**Figure 2** The multiple areas of intervention in our daily consumption in skin surgery that can mitigate impact of dermatological surgery on the climate crisis. PPE, personal protective equipment. Image used with permission from the copyright holders.<sup>1</sup>

However, reusable instruments also exhibit similar perceived attributes. They can be more cost-effective, while also having a lower carbon footprint that is as low as 25% of that of single-use instruments.<sup>8,32</sup> For example, reusable scissors cost £998 with a carbon footprint of 1.87 kg CO<sub>2</sub>e, compared with £24 with 0.475 kg CO<sub>2</sub>e for single use. However, as the reusable device can be used at least 500 times, the overall carbon footprint and costs are lower (£2.00, 0.004 kg CO<sub>2</sub>e per use).<sup>32</sup> Nevertheless, the financial and environmental costs must be weighed between sterilization of reusable instruments, and sharp-bin incineration of single-use instruments.<sup>33</sup> Rizan *et al.* suggested methods to reduce carbon emissions from sterilization, making this a viable solution.<sup>34</sup> Hybrid instruments combining reusable and single-use components can be a cost-effective and environmentally friendly option.<sup>32</sup>

### **Reusing and repurposing used single-use instruments**

Despite instruments being labelled as single use, there are global practices of reusing and reprocessing them for cost savings that raise concerns about patient safety.<sup>35</sup> A systematic review published in 2008 highlighted a lack of direct evidence of the effect of reusing single-use devices on patient outcomes.<sup>35,36</sup>

### **Repairing multiuse equipment**

Repairing and recalibrating reusable instruments when they become ineffective is a sustainable and cost-effective practice, contingent upon suppliers making this option available.<sup>37</sup>

### **Streamlining reusable sets**

Single-centre studies suggest that as much as 87% of instruments in surgical sets remain unused during operations, emphasizing the benefits of streamlining sets with fewer instruments.<sup>38,39</sup> Alternatively, using bespoke instrument sets for different procedures can streamline instrument use and ensure optimal equipment selection.<sup>40,41</sup>

### **Judicious opening of equipment**

Efforts to minimize waste involve judiciously opening equipment when necessary, rather than based on anticipation. A prospective observational single-centre French study showed that more than one-third of surgical waste was caused by supplies prepared in anticipation but not ultimately used.<sup>42</sup> Waste reduction strategies involve waiting until specific phases of procedures to confirm equipment need, fostering better team communication and increasing awareness of supply costs.<sup>42</sup>

### **Sutures**

A meta-analysis of 19 randomized controlled trials (in total, 1748 patients) demonstrated comparable surgical-site infection (SSI) rates and cosmetic outcomes with absorbable vs. nonabsorbable sutures for surface closure.<sup>43</sup> Absorbable sutures reduce plastic waste and obviate the need for suture removal visits.<sup>44</sup> Further research is necessary to explore the environmental impacts associated with the production of different suture materials.

### **Surgical drapes**

Using clean reusable drapes rather than single-use sterile drapes can achieve sufficient infection control. A prospective

study ( $n=1000$ ) evaluated the rate of infection in Mohs micrographic surgery (MMS) using a clean technique (no sterile gloves, gown or drapes) for tumour extirpation and reconstruction. The study found a low incidence of SSIs (0.91%) when using clean (nonsterile) gloves and surgical drapes in MMS. The infection risk for flaps was highest at 2.7% (4 of 150).<sup>45</sup>

### **Mohs micrographic surgery trolley and instruments**

Preserving the instruments on a trolley for each patient throughout Mohs stages and reconstruction ( $n=332$ ) demonstrates low SSI rates (2.1%), reduces instrument changes and associated waste of packaging and drapes, and enhances the overall sustainability of the procedure.<sup>46</sup> This requires strict local protocols to avoid errors or contamination.<sup>47</sup>

### **Local anaesthetic batch preparation**

Batch preparation of a large volume of diluted local anaesthetic can avoid packaging waste of individual syringes, reduce consumables used, and is more time-efficient than individual case preparation. However, utility depends on local context and case volume.<sup>48</sup>

### **Packaging**

#### **Recycling and waste management**

Dermatology departments can significantly enhance sustainability by adopting efficient waste management practices through improving labelling, segregation and educational initiatives.<sup>49</sup> A UK-wide service evaluation reported a mean recycle rate of 16% for nonsharps in skin surgery.<sup>50</sup> However, a 2024 UK national sustainability audit with more included centres found a lower mean recycling rate of 8.6%.<sup>51</sup> Current products, prescriptions and devices have poor and inconsistent recycling labelling, leading to contamination and compromised recycling efforts.<sup>40,49,52</sup> The NHS must leverage its purchasing power to incentivize the industry to adopt appropriate recyclable packaging, consistent labelling and circular economy models.<sup>40</sup>

#### **Bioplastics and plastics**

Globally, most plastic is single use, with 40% of plastics used for packaging and 70% of collected plastics incinerated or sent to landfills.<sup>8</sup> Reducing plastic use in packaging in dermatological surgery will significantly enhance sustainability. Bioplastics are derived from natural resources and have varying degrees of biodegradability.<sup>53</sup> Although bioplastics provide an alternative to conventional plastics, their production still requires significant energy and resources; not all biodegrade efficiently in natural environments, and the transition involves higher costs and changes in supply chain logistics.<sup>53</sup>

#### **Single- vs. double-wrapped surgical equipment**

Studies indicate that single-wrapped sterile packaging carried no greater risk of bacterial contamination than double wrapping,<sup>54</sup> while providing cost savings and environmental benefits.<sup>55</sup> Reusable metal containers are emerging as sustainable alternatives.<sup>34</sup>

## Personal protective equipment

### Caps

Use of reusable surgical cloth caps in dermatological surgery has shown no discernible increase in SSIs.<sup>56</sup> Also, washable cloth caps exhibit less microbiological shedding than disposable bouffant hats.<sup>57</sup> This presents an environmentally favourable option.

### Aprons

In a UK survey of 41 Mohs surgeons, 85% did not use sterile gowns during Mohs surgery for the tumour extirpation stage and 83% for the reconstructive stage of MMS.<sup>58</sup> Use of scrubs or polyethylene aprons had nonsignificant differences in SSIs and substantially lower costs.<sup>59</sup>

### Masks and eye protection

A Cochrane review of 2106 patients found insufficient evidence to support or oppose the use of disposable masks in clean surgery from an SSI perspective.<sup>60,61</sup> Reusable masks offer personal protection against blood splatter and droplet spread and also reduce respiratory particle spread and waste compared with single-use masks.<sup>62</sup>

### Gloves

A systematic review of over 10 000 patients indicated comparable SSI rates between procedures using sterile and non-sterile gloves in cutaneous surgery (including MMS), with an SSI rate of 2.2% in both groups ( $P=0.88$ ). For MMS specifically, nonsterile gloves have a low SSI risk ratio of 1.02 (95% confidence interval 0.78–1.34).<sup>63</sup> The risk of SSI with nonsterile gloves may increase with more complex reconstructions compared with simple excisions, with reported SSI rates varying widely (2.7–14.7%).<sup>45,64,65</sup> Nonsterile gloves are cost-effective (95% cheaper)<sup>66</sup> and are an environmentally sustainable alternative (80% reduction in climate impact).<sup>67</sup>

## Electricity

Operating theatres utilize three to six times more energy than other hospital areas due to ventilation, heating, air conditioning and equipment demands.<sup>68</sup> With up to 40% of theatres being vacant daily, efficiency opportunities exist around occupancy and resource use. Transitioning from halogen to light-emitting diode lighting reduces energy consumption by 49%,<sup>69</sup> and switching off lighting, ventilation, heating/cooling and machinery when not in use can reduce energy waste in dermatological surgery operating theatres.<sup>3</sup>

## Water

A life-cycle assessment of an Australian dermatological surgery department estimated that a typical skin excision procedure consumes 10 L of water (including hand washing and cleaning equipment), equating to 0.07 kg CO<sub>2</sub>e emissions.<sup>4</sup> Switching from soap to alcohol-based hand rubs between surgical cases significantly reduces water use. A Cochrane database meta-analysis found that the method of hand disinfection does not affect the incidence of SSIs.<sup>70</sup> Indeed,

current National Institute for Health and Care Excellence guidelines on the prevention of SSIs recommend that soap and water be used for disinfection at the start of the surgical session, with alcohol gel being sufficient for hand disinfection between cases thereafter (unless hands are visibly soiled).<sup>71</sup>

While 'blue' water consumption (direct use for washing and drinking) constitutes 3.7% of the health sector's water footprint, addressing virtual water (used for equipment and pharmaceutical manufacturing) is also crucial.<sup>19</sup> Implementing streamlined theatre packs and reusing surgical instruments reduce both carbon and water footprints.<sup>40,72,73</sup>

## Paper notes

The NHS produced the Five Year Forward View with the aim of going paperless by adapting, streamlining and reducing the environmental burden of using paper in hospitals.<sup>74</sup> This initiative aims to reduce paper waste and mitigate the environmental impacts of paper production, including deforestation, water consumption and energy use. For instance, producing 1 kg of paper requires two to three times its weight in trees.<sup>75</sup>

## Waste

### Surgical waste

Waste in dermatological theatres falls into clinical and general categories. McGain *et al.* audited six theatres, revealing 45% general waste, 32% clinical waste and 23% recyclable waste, with 40% of the general waste being recyclable.<sup>76</sup> Wernham *et al.* highlighted an annual carbon emission of 26 kg CO<sub>2</sub>e from material waste for one treatment centre, translating to 644 kg CO<sub>2</sub>e across 25 UK centres.<sup>40</sup>

Contaminated or infectious clinical waste requires expensive and energy-intensive autoclave treatment before landfill disposal. Segregating waste at the time of generation and ensuring clear labelling to prevent misclassification of non-hazardous waste are vital for reducing the carbon footprint.<sup>77</sup> A US centre found that addressing the lack of education on waste classification and recycling through a waste management education module yielded a 25% reduction in daily hazardous waste.<sup>78</sup> A carbon footprint analysis by MacNeill *et al.* across three hospitals suggested that refining guidelines to restrict the definition of 'hazardous' can reduce theatre waste further.<sup>68</sup>

It is evident that well-executed recycling programmes within operating rooms significantly mitigate the environmental impact by diverting waste away from landfills.<sup>79</sup> Estimates suggest that up to 90% of theatre waste is non-hazardous and potentially recyclable, including items such as surgical gloves, masks and tubing. Recyclable plastics are often wrongly discarded as general or clinical waste.<sup>80</sup> However, a national audit of 30 UK trusts found that only 57% had recycling bins in their skin surgery rooms.<sup>51</sup> A proposed solution is the incorporation of recycling bins in theatres, improved labelling and education on waste segregation. Also, prepackaged supply kits contribute to waste,

but optimizing them, as seen in Potera's study, can avert a substantial amount of this.<sup>80</sup> Appropriate disposal of sharps is crucial, as glass bottles can increase waste volume.<sup>77</sup>

### Environmentally harmful waste

Incineration of clinical waste contributes to pollution and carcinogenic by-products.<sup>81</sup> Melamed suggested that alternatives like autoclaving or microwaving can be more sustainable, but further research is required to determine the safety and efficacy of this.<sup>82</sup> Formalin, a prevalent fixative for histological specimens, presents a toxic and carcinogenic challenge. Formal handling, storage and disposal protocols for chemicals in MMS are pivotal. Lab atmospheric sampling revealed formaldehyde levels exceeding national standards and identified numerous hazardous or carcinogenic substances.<sup>83</sup> Studies suggest alternatives like sealing tissues under vacuum and specific fixatives such as RCL2 and ethanol, emphasizing the need for ongoing exploration of safer options.<sup>84</sup>

A surgical plume contains over 80 toxic chemicals and poses inhalation health risks, such as respiratory pathologies and potential infections (human papillomavirus, HIV, hepatitis B virus).<sup>85</sup> To address this, healthcare providers should use smoke evacuators and personal protective equipment, and select advanced bipolar electrocautery or ultrasonic devices with lower health risks.<sup>75,86</sup> Waste generated from equipment use, notably in Mohs surgery, has prompted the use of more environmentally sustainable alternatives like hyfrecators over electrosurgical units (including electrocoagulation and electrosection), which has significantly reduced per use waste generation.<sup>87</sup>

### Discussion and conclusions

In part 1 of this review, we discuss the urgent need to enhance environmental sustainability in dermatological surgery, primarily through reducing carbon intensity. This involves analysing the necessary processes required to undertake high-quality skin surgery and evaluating methods to reduce their carbon emissions. Areas identified where dermatological surgery must reduce carbon intensity include procuring goods, transport for staff, patients and materials, and operational resource usage and waste.

Key strategies for adopting low-carbon alternatives include selecting environmentally friendly suppliers, reducing pharmaceutical waste, and optimizing resource use by employing reusable instruments and waste segregation strategies. Additionally, sustainability education among healthcare professionals is vital, incorporating eco-friendly practices into medical training to reduce carbon footprints. Ultimately, achieving meaningful progress requires a system-wide strategy that encompasses reducing carbon intensity, minimizing surgical activity, and incorporating environmental considerations into management, research and innovation.

The second part of this review will explore strategies to reduce surgical activity and future environmental considerations, aiming to integrate sustainability comprehensively into dermatological surgery and effectively mitigate

climate change impacts while maintaining patient care standards.<sup>88</sup>

### Learning points

- The substantial carbon footprint of dermatological surgery demands urgent sustainable practices, particularly as climate change escalates skin cancer rates.
- Opting for sustainable travel choices and educating patients and staff on the environmental impact of the pharmaceutical industry can help reduce the carbon intensity of dermatological surgery.
- Utilizing reusable instruments, judicious opening of equipment, and efficient management of consumables in dermatological surgery have proven cost-effective and environmentally friendly.
- Implementing effective recycling, reducing single-use plastics and ensuring proper waste segregation can minimize environmental impact.
- Water can be saved in operating theatres by using alcohol-based hand rubs.
- Energy consumption in operating theatres can be reduced by adopting energy-efficient lighting, equipment and practices, such as switching off unused devices.

### Acknowledgements

We are grateful to the British Society for Dermatological Surgery Sustainability Subgroup and the British Association of Dermatologists Sustainability Subcommittee for their input on the development of the BSDS Sustainability Guidance 2022.

### Funding sources

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

### Conflicts of interest

The authors declare no conflicts of interest.

### Data availability

No new data were generated or analysed in support of this research.

### Ethics statement

Not applicable.

### Patient consent

Not applicable.

### Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's website.

## References

- Ali F, Nikookam Y, Hunt W *et al.* British Society for Dermatological Surgery Sustainability Guidance 2022. Available at: [https://bsds.org.uk/wp-content/uploads/2022/09/2022-BSDS\\_Sustainability\\_Guidance.pdf](https://bsds.org.uk/wp-content/uploads/2022/09/2022-BSDS_Sustainability_Guidance.pdf) (last accessed 23 October 2024).
- Henriques A, Richardson J, eds. *The Triple Bottom Line: Does it All Add Up?* Abingdon-on-Thames: Routledge, 2013.
- National Health Service. Delivering a net zero NHS. Available at: <https://www.england.nhs.uk/greenernhs/a-net-zero-nhs> (last accessed 23 October 2024).
- Tan E, Lim D. Carbon footprint of dermatologic surgery. *Australas J Dermatol* 2021; **62**:e170–7.
- Levell N. GIRFT. Dermatology. Available at: [https://gettingitright-firsttime.co.uk/medical\\_specialties/dermatology](https://gettingitright-firsttime.co.uk/medical_specialties/dermatology) (last accessed 23 October 2024).
- Cancer Research UK. Statistics by cancer type. Available at: <https://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type> (last accessed 23 October 2024).
- Mieczkowska K, Stringer T, Barbieri JS *et al.* Surveying the attitudes of dermatologists regarding climate change. *Br J Dermatol* 2022; **186**:748–50.
- Ali F, Jabouri H, Wernham AGH, Abbott RA. National dermatological surgery sustainability survey: an evaluation of health-care professional beliefs and practices. *Br J Dermatol* 2023; **188** (Suppl. 4):iv113.
- Centre for Sustainable Healthcare. 10 Year Impact Review. Available at: <https://sustainablehealthcare.org.uk/file/1141> (last accessed 23 October 2024).
- Tomson C. Reducing the carbon footprint of hospital-based care. *Future Hosp J* 2015; **2**:57–62.
- Niebel D, Herrmann A, Balzer S *et al.* Sustainability of dermatological offices and clinics: challenges and potential solutions. *J Dtsch Dermatol Ges* 2023; **21**:44–58.
- Altman DJ. The roles of pharmaceutical industry and drug development in dermatology and dermatologic health care. *Dermatol Clin* 2000; **18**:287–96.
- Ooi K. The pitfalls of overtreatment: why more care is not necessarily beneficial. *Asian Bioeth Rev* 2020; **12**:399.
- Academy of Medical Royal Colleges. Protecting resources, promoting value: a doctor's guide to cutting waste in clinical care. Available at: <https://www.hfma.org.uk/publications/protecting-resources-promoting-value-doctors-guide-cutting-waste-clinical-care-academy> (last accessed 23 October 2024).
- Pawar P, Reddy S, Bharat Gaikwad D *et al.* Reducing waste to create a sustainable supply and usage of pharmaceuticals. *Int J Adv Res Sci Commun Technol* 2023; **3**:297–302.
- BASF. It's time to calculate the carbon footprint of pharmaceutical products. Available at: <https://themedicinemaker.com/manufacture/its-time-to-calculate-the-carbon-footprint-of-pharmaceutical-products> (last accessed 23 October 2024).
- Llano Martinez G. Environmental impact of the pharmaceutical packaging. Available at: <https://lup.lub.lu.se/student-papers/search/publication/3044827> (last accessed 23 October 2024).
- Fathy R, Nelson CA, Barbieri JS. Combating climate change in the clinic: cost-effective strategies to decrease the carbon footprint of outpatient dermatologic practice. *Int J Womens Dermatol* 2021; **7**:107–11.
- NHS England. Reducing the use of natural resources in health and social care, 2018 report. Available at: <https://healthacademy.lancsteachinghospitals.nhs.uk/app/uploads/2022/02/Reducing-the-use-of-natural-resources-in-health-and-social-care.pdf> (last accessed 23 October 2024).
- Watts N, Amann M, Arnell N *et al.* The 2019 report of The Lancet Countdown on health and climate change: ensuring that the health of a child born today is not defined by a changing climate. *Lancet* 2019; **394**:1836–78.
- NHS England. Early diagnosis of skin cancer: innovating the two-week wait skin cancer referral pathway as part of the NHS post COVID-19 targeted intervention recovery plan. Available at: <https://cdn.bad.org.uk/uploads/2022/02/29200037/Suspected-Skin-Cancer-2ww-Pathway-Optimisation-Proposal-v19-FINAL-16.06.2021.pdf> (last accessed 23 October 2024).
- NESTA. Transforming elective care services: dermatology. Available at: <https://www.england.nhs.uk/wp-content/uploads/2019/01/dermatology-elective-care-handbook-v1.pdf> (last accessed 23 October 2024).
- General Medical Council. Outcomes for graduates, plus supplementary guidance. Available at: <https://www.gmc-uk.org/education/standards-guidance-and-curricula/standards-and-outcomes/outcomes-for-graduates> (last accessed 23 October 2024).
- Rizan C, Reed M, Mortimer F *et al.* Using surgical sustainability principles to improve planetary health and optimise surgical services following the COVID-19 pandemic. *Bull R Coll Surg Engl* 2020; **102**:177–81.
- Chanchlani A, Martin K, Dunne B. Environmental sustainability in simulation education in healthcare. *Int J Healthc Simul* 2024; **10**:54531/mqqv2910.
- Laspro M, Groysman L, Verzella A *et al.* The use of virtual reality in surgical training: implications for education, patient safety, and global health equity. *Surgeries* 2023; **4**:635–46.
- Ha ES, Hong JY, Lim SS *et al.* The impact of SARS-CoV-2 (COVID-19) pandemic on international dermatology conferences in 2020. *Front Med* 2021; **8**:1266.
- Privalle A, Havighurst T, Kim KM *et al.* Number of skin biopsies needed per malignancy: comparing the use of skin biopsies among dermatologists and nondermatologist clinicians. *J Am Acad Dermatol* 2020; **82**:110–16.
- Thiel CL, Eckelman M, Guido R *et al.* Environmental impacts of surgical procedures: life cycle assessment of hysterectomy in the United States. *Environ Sci Technol* 2015; **49**:1779.
- Drew J, Christie SD, Rainham D, Rizan C. HealthcareLCA: an open-access living database of health-care environmental impact assessments. *Lancet Planet Health* 2022; **6**:e1000–12.
- Parkins GJ, Wylie G. Dermatological surgery – time for single-use instruments? *Dermatol Surg* 2014; **40**:1434.
- Rizan C, Bhutta MF. Environmental impact and life cycle financial cost of hybrid (reusable/single-use) instruments versus single-use equivalents in laparoscopic cholecystectomy. *Surg Endosc* 2021; **36**:4067–78.
- Basu D, Dutta SK, Bhattacharya S *et al.* The economics of autoclave-based sterilization: experience from central sterile supply department of a cancer center in Eastern India. *Infect Control Hosp Epidemiol* 2016; **37**:878–80.
- Rizan C, Lillywhite R, Reed M, Bhutta MF. Minimizing carbon and financial costs of steam sterilization and packaging of reusable surgical instruments. *Br J Surg* 2021; **109**:200–10.
- Hailey D, Jacobs PD, Ries NM, Polisena J. Reuse of single use medical devices in Canada: clinical and economic outcomes, legal and ethical issues, and current hospital practice. *Int J Technol Assess Health Care* 2008; **24**:430–6.
- Jacobs P, Polisena J, Hailey D, Lafferty S. Economic analysis of reprocessing single-use medical devices: a systematic literature review. *Infect Control Hosp Epidemiol* 2008; **29**:297–301.
- Medicines and Healthcare products Regulatory Agency. Managing medical devices. Available at: <https://www.gov.uk/government/publications/managing-medical-devices> (last accessed 23 October 2024).
- Stockert EW, Langerman A. Assessing the magnitude and costs of intraoperative inefficiencies attributable to surgical instrument trays. *J Am Coll Surg* 2014; **219**:646–55.
- Nast K, Swords KA. Decreasing operating room costs via reduction of surgical instruments. *J Pediatr Urol* 2019; **15**:153.
- Wernham A, Patel A, Sharma A. Environmental impact of Mohs surgery and measures to reduce our carbon footprint in dermatological surgery. *Br J Dermatol* 2019; **181** (Suppl. S1):105.

- 41 Cichos KH, Hyde ZB, Mabry SE *et al.* Optimization of orthopedic surgical instrument trays: lean principles to reduce fixed operating room expenses. *J Arthroplasty* 2019; **34**:2834–40.
- 42 Chasseigne V, Leguelinel-Blache G, Nguyen TL *et al.* Assessing the costs of disposable and reusable supplies wasted during surgeries. *Int J Surg* 2018; **53**:18–23.
- 43 Xu B, Xu B, Wang L *et al.* Absorbable versus nonabsorbable sutures for skin closure: a meta-analysis of randomized controlled trials. *Ann Plast Surg* 2016; **76**:598–606.
- 44 Parell GJ, Becker GD. Comparison of absorbable with nonabsorbable sutures in closure of facial skin wounds. *Arch Facial Plast Surg* 2003; **5**:488–90.
- 45 Rogers HD, Desciak EB, Marcus RP *et al.* Prospective study of wound infections in Mohs micrographic surgery using clean surgical technique in the absence of prophylactic antibiotics. *J Am Acad Dermatol* 2010; **63**:842–51.
- 46 Nasser E. Prospective study of wound infections in Mohs micrographic surgery using a single set of instruments. *Dermatol Surg* 2015; **41**:1008–12.
- 47 Petukhova TA, King TH, Omlin KJ, Eisem DB. Reusing surgical instruments during Mohs micrographic surgery: safe from infection, but not free from risk. *Dermatol Online J* 2016; **22**:13030/qt3430b8nw.
- 48 Gleeson C, Hussain W, Spreadborough J *et al.* Local anaesthetic preparation in dermatological surgery: a labour- and time-efficient approach. *Br J Dermatol* 2011; **164**:888–90.
- 49 Tso VBY, Lambreghts CS, Tso S *et al.* On-pack recycling label in cosmeceutical products in dermatology. *Clin Exp Dermatol* 2022; **47**:186–8.
- 50 Shearman H, Yap SM, Zhao A *et al.* A UK-wide study to describe resource consumption and waste management practices in skin surgery including Mohs micrographic surgery. *Clin Exp Dermatol* 2023; **48**:1024–9.
- 51 Forbat E, Barlow R, Wernham A *et al.* The British Society for Dermatological Surgery's national sustainable skin surgery audit standards and checklist: preliminary findings and recommendations from the first national audit. *Br J Dermatol* 2024; **191** (Suppl. 1):i95.
- 52 Kamp E, Musbahi E, Gupta P. Topical treatment samples: plastic, recycling and sustainability. *Clin Exp Dermatol* 2022; **47**:186.
- 53 Narancic T, Cerrone F, Beagan N, O'Connor KE. Recent advances in bioplastics: application and biodegradation. *Polymers (Basel)* 2020; **12**:920.
- 54 Webster J, Radke E, George N *et al.* Barrier properties and cost implications of a single versus a double wrap for storing sterile instrument packs. *Am J Infect Control* 2005; **33**:348–52.
- 55 Krohn M, Fengler J, Mickley T, Flessa S. Analysis of processes and costs of alternative packaging options of sterile goods in hospitals – a case study in two German hospitals. *Health Econ Rev* 2019; **9**:1.
- 56 ARHAI Scotland. PPE Headwear literature review. Available at: <https://www.nipcm.hps.scot.nhs.uk/media/1669/2021-08-18-sicp-lr-headwear-v3.pdf> (last accessed 23 October 2024).
- 57 Markel TA, Gormley T, Greeley D *et al.* Hats off: a study of different operating room headgear assessed by environmental quality indicators. *J Am Coll Surg* 2017; **225**:573–81.
- 58 Chan BCY, Al-Niaimi F, Mallipeddi R. Infection control practices in Mohs micrographic surgery: a U.K. national survey. *Br J Dermatol* 2017; **177**:e204–5.
- 59 Lilly E, Schmults CD. A comparison of high- and low-cost infection-control practices in dermatologic surgery. *Arch Dermatol* 2012; **148**:859–61.
- 60 Kupres K, Rasmussen SE, Albertini JG. Perforation rates for non-sterile examination gloves in routine dermatologic procedures. *Dermatol Surg* 2002; **28**:388–9.
- 61 Vincent M, Edwards P. Disposable surgical face masks for preventing surgical wound infection in clean surgery. *Cochrane Database Syst Rev* 2016; **4**:CD002929.
- 62 Asadi S, Cappa CD, Barreda S *et al.* Efficacy of masks and face coverings in controlling outward aerosol particle emission from expiratory activities. *Sci Rep* 2020; **10**:15665.
- 63 Brewer JD, Gonzalez AB, Baum CL *et al.* Comparison of sterile versus nonsterile gloves in cutaneous surgery and common outpatient dental procedures: a systematic review and meta-analysis. *JAMA Dermatol* 2016; **152**:1008–14.
- 64 Rogues AM, Lasheras A, Amici JM *et al.* Infection control practices and infectious complications in dermatological surgery. *J Hosp Infect* 2007; **65**:258–63.
- 65 Schwartzman G, Khachemoune A. Surgical site infection after dermatologic procedures: critical reassessment of risk factors and reappraisal of rates and causes. *Am J Clin Dermatol* 2021; **22**:503–10.
- 66 Riyaz FR, Ozog DM, Dover JS. Managing and reducing office expenses in dermatology surgery. *Dermatol Surg* 2020; **46**:443–5.
- 67 Jamal H, Lyne A, Ashley P, Duane B. Non-sterile examination gloves and sterile surgical gloves: which are more sustainable? *J Hosp Infect* 2021; **118**:87–95.
- 68 MacNeill AJ, Lillywhite R, Brown CJ. The impact of surgery on global climate: a carbon footprinting study of operating theatres in three health systems. *Lancet Planet Heal* 2017; **1**:e360–7.
- 69 Stall NM, Kagoma YK, Bondy JN, Naudie D. Surgical waste audit of 5 total knee arthroplasties. *Can J Surg* 2013; **56**:97.
- 70 Tanner J, Dumville JC, Norman G, Fortnam M. Surgical hand antisepsis to reduce surgical site infection. *Cochrane Database Syst Rev* 2016; **2016**:CD004288.
- 71 National Institute for Health and Care Excellence. Surgical site infections: prevention and treatment. Available at: <https://www.nice.org.uk/guidance/ng125/chapter/recommendations> (last accessed 23 October 2024).
- 72 Van Demark RE, Smith VJS, Fiegen A. Lean and green hand surgery. *J Hand Surg Am* 2018; **43**:179–81.
- 73 NHS Sustainable Development Unit. Case study. Waste. Royal Liverpool & Broadgreen University Hospitals. Available at: [https://www.molnlycke.com/globalassets/royal\\_liverpool\\_broadgreen—environmental-study.pdf](https://www.molnlycke.com/globalassets/royal_liverpool_broadgreen—environmental-study.pdf) (last accessed 23 October 2024).
- 74 NHS. Next steps on the NHS Five Year Forward View. Available at: <https://www.england.nhs.uk/wp-content/uploads/2017/03/NEXT-STEPS-ON-THE-NHS-FIVE-YEAR-FORWARD-VIEW.pdf> (last accessed 23 October 2024).
- 75 Lewin JM, Brauer JA, Ostad A. Surgical smoke and the dermatologist. *J Am Acad Dermatol* 2011; **65**:636–41.
- 76 McGain F, Jarosz KM, Nguyen MNHH *et al.* Auditing operating room recycling: a management case report. *A A Case Rep* 2015; **5**:47–50.
- 77 Wyssusek KH, Foong WM, Steel C, Gillespie BM. The gold in garbage: implementing a waste segregation and recycling initiative. *AORN J* 2016; **103**:316.
- 78 Samaan C, Kamrani P, Ken KM. Reducing environmental impact in dermatology: a single-institution educational approach to reducing biohazard waste. *Arch Dermatol Res* 2023; **315**:2755–6.
- 79 de'Angelis N, Conso C, Bianchi G *et al.* Systematic review of carbon footprint of surgical procedures. *J Visc Surg* 2024; **161** (2 Suppl.):7–14.
- 80 Potera C. Strategies for greener hospital operating rooms. *Environ Health Perspect* 2012; **120**:a306.
- 81 Wyssusek KH, Keys MT, van Zundert AAJ. Operating room greening initiatives – the old, the new, and the way forward: a narrative review. *Waste Manag Res* 2019; **37**:3–19.
- 82 Melamed A. Environmental accountability in perioperative settings. *AORN J* 2003; **77**:1157–68.
- 83 Gunson TH, Smith HR, Vinciullo C. Assessment and management of chemical exposure in the Mohs laboratory. *Dermatol Surg* 2011; **37**:1–9.
- 84 Sarot E, Carillo-Baraglioli MF, Duranthon F *et al.* Assessment of alternatives to environmental toxic formalin for DNA conservation in biological specimens. *Environ Sci Pollut Res Int* 2017; **24**:16985–93.



- 85 Ball K. Update for nurse anesthetists. Part 1. The hazards of surgical smoke. *AANA J* 2001; **69**:125–32.
- 86 Ball K. Surgical smoke evacuation guidelines: compliance among perioperative nurses. *AORN J* 2010; **92**:e1–23.
- 87 Leonard N, McLean-Mandell R. A step toward environmental sustainability in Mohs surgery. *Dermatol Surg* 2021; **47**:1504–5.
- 88 Ali F, Wernham A, Abbott R. Environmental sustainability in dermatological surgery. Part 2: reducing activity and future ecological strategies. *Clin Exp Dermatol* 2025; in press

## Appendix 1

### BSDS Sustainability Subgroup Collaborative

The BSDS Sustainability Subgroup Collaborative consists of Fatima Ali, Rachel Abbott, Aaron Wernham, Yasmin Nikookam, William Hunt, Sophie Holloran, Catriona Chaolin, Eshen Ang, Maria Charalambides, Ashima Lowe, Luke Brindley, Christopher Bower, Sandeep Varma, Minh Lam, David Veitch, Hilmi Recica, Wen Ai Woo, Simon Tso and Claire Doyle.

### CPD questions

#### Learning objective

To become more familiar with environmental sustainability in dermatological surgery.

#### Question 1

Which category contributes the most to the National Health Service's carbon footprint?

- (a) Medical equipment.
- (b) Paper notes.
- (c) Patient travel.
- (d) Staff travel.
- (e) Supply chain and procurement.

#### Question 2

Which of the following is an effective strategy to minimize water use in dermatological surgery?

- (a) Adopting alcohol-based hand rubs between cases.

- (b) Increasing the water temperature for hand washing.
- (c) Not washing hands between cases.
- (d) Using disposable hand towels.
- (e) Using soap and water for all hand disinfections.

#### Question 3

How can dermatological surgery departments reduce their electricity consumption?

- (a) Always keeping operating theatre lights and equipment on.
- (b) Extending operating theatre hours.
- (c) Implementing energy-efficient light-emitting diode lighting and turning off unused equipment.
- (d) Installing additional air conditioning units in rooms.
- (e) Using halogen lighting in operating theatres.

#### Question 4

Which of the following practices helps in reducing waste in dermatological surgery?

- (a) Disposing of unused sterile equipment after each surgery.
- (b) Enhancing recycling efforts and proper waste segregation.
- (c) Ignoring the segregation of waste types.
- (d) Increasing use of double-wrapped instruments.
- (e) Using more single-use instruments instead of reusable ones.

#### Question 5

What are considered sustainable travel options that reduce carbon emissions of healthcare?

- (a) Encouraging the use of diesel vehicles for all travel.
- (b) Increasing the use of cars for patient and staff transport.
- (c) Allowing staff and patient transport only using hybrid vehicles.
- (d) Promoting the use of cycling, walking and public transport.
- (e) Relying solely on traditional taxis for patient and staff transport.